

Genetic diversity of the novel coronavirus SARS-CoV-2 (COVID-19) in Portugal

More information at <https://insaflu.insa.pt/covid19>



Situation Report

December 27th, 2022

The National Institute of Health Doutor Ricardo Jorge, I.P. (INSA) has analysed **44385** SARS-CoV-2 genome sequences so far.




Figure 1: Evolution of the weekly relative frequency of the SARS-CoV-2 variants circulating in Portugal between ISO weeks 22 (31/05/21 - 06/06/21) and 50 (13/12/22 - 19/12/22). The frequencies presented for the last week under analysis (ISO week 50*) might change in the next report, given that some data from that period is still being processed. *This and other graphs can be explored interactively on the website.*

Main highlights

- **Lineage BA.5 of the variant Omicron** (including its multiple sub-lineages) **is dominant in Portugal** since week 19 (09/05/22 - 15/05/22) and presents a **relative frequency of 94,6%** according to the most recent national sequencing survey on week 50 (13/12/22 - 19/12/22) (**Figure 1**).
- The relative frequency of **lineage BA.4 of the variant Omicron** has been residual, with **no sequence detected in the last 2 weeks** (**Figure 1**).
- **Lineage BA.2 of the variant Omicron was dominant in Portugal between weeks 8 (21/02/22 - 27/02/22) and 19 (09/05/22 - 15/05/22)**. Since then, its relative frequency is residual, being represented mainly by **lineage BN.1 (and its sub-lineages)**, with relative frequencies of 3,7% and 3,6% in weeks 49 and 50, respectively.
- On behalf of the continuous monitoring of the introduction and circulation of (new) SARS-CoV-2 (sub-)lineages in Portugal, **we have been observing the emergence of sub-lineages of interest**, with novel constellations of mutations potentially associated with resistance to neutralizing antibodies. Among these, we highlight the **sub-lineages of BA.2** (e.g., BN.1 and CA.7) **and BA.5** (e.g., BF.7 and BQ.1), some of them showing considerable increase of circulation in some countries, particularly in Europe. In Portugal, it is highlighted the **increase of the relative frequencies of BQ.1** (and its sub-lineages, in particular BQ.1.1), **which is dominant since week 44** (**Figure 2**). To date, 29 sequences of the recombinant sub-lineage XBB have been detected in Portugal. XBB has been a focus of interest due to its immune evasion capacity.




Figure 2: Evolution of the relative frequency of sub-lineage BQ.1 and its descendants BQ.1.1 and BQ.1.1.15 in Portugal. The circular graph shows the distribution of the relative frequencies of SARS-CoV-2 sub-lineages in the period of ISO weeks 49 and 50 (06/12/22 - 19/12/22), **highlighting sub-lineages representing >5% of the sequences analysed in this period**. The evolution of BQ.1** and BQ.1.1** relative frequencies during the last 14 weeks is shown in the line plot. It is expected that the frequencies presented for the last week under analysis (ISO week 50*) might change in the next report, given that some data from that period is still being processed. **the presented relative frequencies correspond to the sub-lineages and their descendants. *Other graphs can be explored interactively on the website.*

Autorship

Genomics and Bioinformatics Unit
Department of Infectious Diseases
NATIONAL INSTITUTE OF HEALTH DOUTOR RICARDO JORGE
Avenida Padre Cruz, 1649-016 Lisboa, PORTUGAL

Recommended citation

National Institute of Health (INSA) Doutor Ricardo Jorge. Genetic diversity of the novel coronavirus SARS-CoV-2 (COVID-19) in Portugal. Lisbon, Portugal INSA; 2022. Available at: <https://insaflu.insa.pt/covid19>

Useful links

<https://www.ecdc.europa.eu/en/covid-19/situation-updates/variants-dashboard>
<https://www.who.int/activities/tracking-sars-cov-2-variants>
https://cov-lineages.org/lineage_list.html
<https://outbreak.info/>
<https://www.gisaid.org/>